

State of the Primary Care Workforce, 2024

November 2024

Primary care is a fundamental part of the nation's health care system. Better access to and use of primary care has been shown to improve treatment of chronic conditions and increase life expectancy. However, it is well-documented that significant challenges face the workforce providing this care.¹ These include shortages and maldistribution of primary care providers (PCPs), low compensation compared to other health occupations, increasing burnout and job dissatisfaction, and an aging workforce.

The primary care workforce is defined in this report as physicians, nurse practitioners (NPs), and physician assistants (PAs) practicing in primary care specialties: family medicine, general pediatric medicine, general internal medicine, and geriatric medicine. While the majority of the nation's hospitalists—providers who mainly provide care to hospitalized patients—are trained in primary care specialties, they are excluded from provider counts in this report (except where noted) as these clinicians are not engaged in activities that meet the definition of primary care.² All physicians, NPs, and PAs in this report refer to the primary care workforce unless noted.

The purpose of this report is to update and discuss HRSA's most recent projected estimates of the future supply of primary care occupations and give context for that workforce by examining their current state.

Highlights

- In 2022, there were 279,194 primary care physicians in the U.S. In 2022, there were an estimated 270,660 NPs delivering primary care. In 2023, 28,282 PAs worked in primary care.
- There is a projected shortage of 87,150 full-time equivalent (FTE) primary care physicians by 2037, which will be particularly acute in nonmetro areas.
- A substantial and increasing amount of behavioral health and obstetrics and gynecology (OB-GYN) services are being provided by PCPs.
- Primary care physicians, NPs, and PAs earn less than counterparts in other specialties.
- Burnout has increased in many healthcare occupations, but especially among primary care physicians. More than half reported feeling burnout in 2022.
- Primary care physicians used telehealth more during the COVID-19 pandemic than prior to 2020.
- The demographics and geographic location of the U.S population are projected to change dramatically over the next 35 years. The primary care workforce will have to change with it to continue to deliver high-quality care.

About the National Center for Health Workforce Analysis

The National Center for Health Workforce Analysis informs public and private sector decision makers on health workforce issues by expanding and improving health workforce data, disseminating workforce data to the public, and improving and updating projections of the supply and demand for health workers.

For more information, visit the [Health Workforce Analysis](#) webpage.

Describing the primary care workforce

PCPs are often the first point of contact for patients seeking medical care. They play a vital role in preventive care, early detection and treatment of diseases, management of chronic conditions, and acute care in both inpatient and outpatient settings.^{3,4} PCPs also play a crucial role in the provision of behavioral health and women’s health services.^{5,6} Overall, this workforce is vital for the U.S. population to remain healthy, manage diseases, and prevent illnesses and deaths.⁷

Enumeration

In 2022, 279,194 primary care physicians were actively working, representing 29.9% of all U.S. active physicians.⁸ From 2016 to 2021, the number of primary care physicians increased by only 3.6%, in contrast to an 8.7% increase among other (including unknown) specialties.⁹ These figures include only primary care physicians who met these criteria: actively practicing, younger than age 75, and not in residency. Table 1 presents the breakdown of these physicians by specialty.

Table 1. Active Primary Care Physicians by Specialty in 2022, Number and Percent

Physician Specialty	Count	Percentage
Family medicine	107,536	38.5%
Internists	106,294	38.1%
Geriatricians	5,611	2.0%
Pediatricians	59,753	21.4%
All primary care physicians	279,194	100%

Note. 2022 AMA Physician Professional Data.

One striking finding is the low number and percentage of geriatricians in the U.S. This is notable due to the well-documented large number of U.S. adults aged 65 and older, who might require the care of a geriatrician. Additionally, an estimated 270,660 NPs in 2022 and 28,282 PAs in 2023 provided primary care.^{10,11} PAs with a secondary position in primary care were not included in this estimate. While NPs and PAs do not have the same level of training and autonomy as primary care physicians, they do deliver primary care services.

Demographics

The primary care physician workforce varies demographically depending on the specialty (Table 2).

Table 2. Demographics of Primary Care Physicians by Specialty Type

Demographics	Family Medicine	Internists	Geriatricians	Pediatricians
Men ^a	54.9%	58.5%	43.3%	32.7%
Women ^a	45.1%	41.5%	56.7%	67.3%
Ages 34 and younger ^b	8.1%	10.0%	6.2%	9.6%
Ages 35 to 44 ^b	22.7%	22.8%	27.0%	23.8%
Ages 45 to 54 ^b	27.0%	25.3%	33.0%	28.7%
Ages 55 to 64 ^b	24.4%	25.5%	21.1%	23.6%
Ages 65 and older ^b	17.8%	16.3%	12.7%	14.3%
White ^c	66.3%	52.7%	44.1%	64.5%
Black/African American ^c	6.7%	7.8%	7.2%	7.4%
Asian ^c	16.2%	29.4%	35.4%	17.0%
Other ^c	2.9%	2.9%	3.9%	2.4%
Hispanic or Latino	7.9%	7.2%	9.4%	8.7%

Note. Sex and age data are from the 2022 AMA Physician Professional Data and exclude hospitalists; other data in the table are from the AAMC Physician Specialty Data Report prepared based on analysis of AMA Physician Professional Data (Dec. 31, 2021) and include hospitalists.

^a Excludes unknown sex. ^b Excludes unknown age. ^c Non-Hispanic or Latino.


Women make up most geriatricians and pediatricians while there are more male family medicine physicians and internists. Further, over 40% of family medicine physicians and internists are age 55 and older.

For all NPs (not just primary care), 90% are women, 78% are non-Hispanic White, and the median age is 42.7.¹² PAs (not just primary care) are predominately non-Hispanic White (72%), 39 years and younger (59%), and female (66%).¹³

Distribution

The distribution of primary care physicians differs by level of urbanization. In general, rural areas have lower primary care physician ratios than urban areas.^{14,15,16} In 2022, 7.8% of U.S. counties did not have a primary care physician at all and the national ratio of primary care physicians was 83.8 per 100,000 population.^{8,17} Whether or not this is considered adequate at the national level, the range of ratios across the states shows an uneven distribution of these physicians (*Figure 1 and Table A in Appendix*). Often a national maldistribution is interpreted as a shortage at the state (or lower) geographic level.

Figure 1. Ratios of Primary Care Physicians per 100,000 Population by U.S. State, 2022



Note. 2022 AMA Physician Professional Data and the U.S. Census Bureau's State Population Totals: 2020-2023 (census.gov).

NPs and PAs are important in providing primary care in rural areas. Approximately half of PAs were interested in practicing in rural locations (44%), Medically Underserved Areas (MUAs) (58%), or Health Professional Shortage Areas (HPSAs) (54%).^{18,19}

Current and projected shortages

As of June 30, 2024, there are 7,501 designated primary care HPSAs in the United States, with nearly 75 million residents (approximately 22% of the U.S. population).²⁰ According to the most recent data, 66.5% of designated primary care HPSAs are in rural areas. Based on a minimum adequate population-to-primary care physician ratio of 3,500 to 1, HRSA estimates that the United States needs 13,075 additional physicians to remove all primary care shortage designations.

As for the future, HRSA projects a national shortage of 87,150 full-time equivalent (FTE) primary care physicians by the year 2037.²¹ To determine if the number of physicians in a specialty will be adequate, the projected supply is subtracted from the projected demand. As seen in Table 3, all primary care physician specialties will experience some level of shortage ranging from 73% adequacy (family medicine physicians) to 81% adequacy (pediatricians) in 2037. The projected supply of family medicine physicians in 2037 will be sufficient to meet only 73% of demand in that year; stated simply, there will be a 27% shortage of these physicians. There are also significant differences in projected shortages between metro and nonmetro areas.²¹

Table 3. Projected Shortage of Primary Care Physicians by Specialty in 2037, Number and Percent Adequacy

Physician Specialty	Metro	Nonmetro	All United States
Family medicine	35,910 (74%)	7,310 (68%)	43,220 (73%)
Geriatricians	1,560 (82%)	550 (34%)	2,110 (78%)
Internists	21,990 (80%)	6,900 (42%)	28,890 (76%)
Pediatricians	9,940 (84%)	2,990 (53%)	12,930 (81%)
Total	69,400 (78%)	17,750 (58%)	87,150 (76%)

Note. HRSA Workforce Projections - <https://data.hrsa.gov/topics/health-workforce/workforce-projections>.

A major factor contributing to the projected shortage of primary care physicians in the future is the age of primary care physicians. The primary care physician workforce is older than other occupations, which means higher rates will be leaving the labor force in the coming decades.

NPs and PAs may to some degree alleviate the issues associated with the shortage of primary care physicians. There is a projected surplus of nurse practitioners (66,360 FTEs) and small projected surplus of PAs (460 FTEs) in 2037.

Challenges for the Primary Care Workforce

Compensation

One of the main challenges to attract new clinicians to the primary care workforce is low compensation relative to other clinicians. Primary care is among the lowest paid physician fields. Table 4 shows 2023 annual average salaries for the selected physician specialties.

Table 4. Earnings for Selected Physician Specialties, 2023

Physician Specialty	Annual Salary
Orthopedics	\$558,000
Plastic surgery	\$536,000
General surgery	\$423,000
Obstetrics and gynecology	\$352,000
Psychiatry	\$323,000
<i>Internal medicine^a</i>	\$282,000
<i>Family medicine^a</i>	\$272,000
<i>Pediatrics^a</i>	\$260,000

Note. 2024 Medscape's Physician Compensation Report (geriatrics was not included in the report). ^a Indicates primary care specialties.

The substantial gap in compensation between primary care physicians and specialist physicians may be one of the explanatory factors for medical students choosing residency in specialties other than primary care.²² Further, salaries for NPs and PAs working in primary care are lower than the average salaries of their counterparts outside of primary care. The average NP salary in 2021 was \$113,000, and a reported average salary for NPs working in primary care was \$100,820 in 2022.^{10,23} The 2023 median salary for all PAs was \$130,020; PAs working in primary care earned a median annual salary of \$115,000 in 2023.^{11,24}

Burnout

Primary care physicians have high rates of burnout, with burnout increasing in recent years.^{25,26} Recent studies show that rates of burnout for physicians increased from 42% in 2020 to 47% and 53% in 2021 and 2022, respectively, before declining slightly to 49% in 2023.²⁷ Compared to other physician specialties, two of the four primary care specialties are among the five specialties reporting the highest level of burnout (Table 5).

Table 5. Percentage of Physicians Reporting Burnout in Selected Specialties, 2023

Physician Specialty	Percent of Physicians who Reported Burnout
Emergency medicine	63%
Obstetrics and gynecology	53%
Oncology	53%
<i>Pediatrics^a</i>	51%
<i>Family medicine^a</i>	51%
<i>Internal medicine^a</i>	50%
General surgery	45%
Psychiatry	39%

Note. <https://www.medscape.com/slideshow/2024-lifestyle-burnout-6016865#3>. ^a Indicates primary care specialties.

High burnout rates and low satisfaction with work-life balance has been consistently reported by primary care physicians in the past decade. Studies that surveyed U.S. physicians in 2011, 2014, 2017, and 2020 found family medicine physicians and internists had high burnout rates and low satisfaction with work-life integration.^{28,29} Primary care physicians reported higher rates of burnout in the 2021 survey when compared to the surveys conducted in 2011, 2014, 2017 and 2020.³⁰

Common factors contributing to physician burnout are workload, long working hours, clerical duties, and a large number of patients.³¹ In addition, violence in the workplace contributes negatively to health care workforce wellbeing.^{29,32} The health care workforce experiences higher rates of workplace violence than workers in other industries.^{33,34}

Telehealth

Since 2020, health care providers increased their use of telehealth resources.³⁵ A recent study evaluating the use of telehealth before and during the COVID-19 pandemic found that only 5.3% of primary care physicians used telehealth “often” before the COVID-19 pandemic, while nearly half (46.2%) reported using telehealth often during the COVID-19 pandemic (*Table 6*).³⁶ Telehealth increases access to care in shortage areas, reduces travel and wait time for patients, and increases access for patients with limited mobility as well as collaboration between care providers.^{37,38} Telehealth proved to be a highly effective instrument in connecting health care providers and patients during the pandemic.

Table 6. Telehealth Use by Primary Care Physicians

Telehealth Use	Pre COVID-19	During COVID-19	Intent to Use After COVID-19
Often	5.3%	46.2%	26.2%
Occasionally	13.4%	34.4%	46.6%
Rarely	24.9%	11.7%	16.3%
Never	54.6%	5.9%	9.1%
Missing (not reported)	1.8%	1.8%	1.8%

Note. Adapted from Callaghan T., McCord C., Washburn D., Goidel K., Schmit C., Nuzhath T., Spiegelman A., & Scobee J. (2022). The Changing Nature of Telehealth Use by Primary Care Physicians in the United States. *Journal of Primary Care & Community Health*. 13:1-9. <https://doi.org/10.1177/21501319221110418>.

Population factors impacting the primary care workforce

The U.S. population grew rapidly from 1980 to 2020, increasing 46% (227 million to 331 million).³⁹ The future U.S. population is predicted to increase 10% (331 million to 364 million) from 2020 to 2060.⁴⁰ The number of primary care physicians is increasing at approximately the same rate as the population they serve. It is estimated that in 2037, the national ratio of FTE primary care physicians will be 76.8 per 100,000 individuals as compared to 81.6 in 2022.^{21,40}

The 2020 Census revealed significant geographic shifts in the United States population. Among the 2.1 million U.S. residents who changed their region of residence in 2022, 63.8% moved to the South and West regions. This will have a potentially huge impact on the distribution needs for the primary care workforce now and into the future.

Demographics

The 65 and older population is projected to increase 54% (from 58 million to 89 million) between 2022 and 2060, with nearly 1 in 4 Americans being 65 years and older in 2060.⁴¹ This trend will have significant implications for the health care industry as the demand for the services related to an older population will surge.⁴²

Between 2022 and 2060, the population of women in the United States is projected to grow nearly 10% (from approximately 168 million to 184 million).⁴³ Since many primary care physicians provide women's health services, this growth will add to the future demand for primary care physicians.^{44, 45, 46}

By 2060, the percentage of the population in demographic groups other than non-Hispanic White will increase from 41.1% in 2022 to 55.1% in 2060.⁴⁷

Improving population health via the primary care workforce

Access to and Use of Primary Care Providers

Health care access is generally defined as the ability to obtain health care services in a convenient and affordable way. Studies have shown that better access to primary care providers leads to improved health outcomes for the population.^{7,48} Barriers to accessing primary care providers in the United States include shortages and geographical maldistribution of providers, transportation issues, lack of health insurance, and limited office hours.^{49,50}

The percentage of the U.S. population having a usual source of care has declined in recent years.⁵¹ The usual source of care is a medical professional or facility where an individual regularly accesses medical care and is typically a primary care provider.^{52,53} This is partially why behavioral health care and OB-GYN services are becoming an increasingly large part of primary care visits.⁵⁴ A recent study found that the share of primary care visits that addressed mental and behavioral concerns increased by 49% from the period between 2006-2007 and 2016-2018.⁵⁵ Primary care physicians often screen patients during their primary care visits for behavioral health issues and prescribe and manage medications to treat depression, substance abuse, and attention deficit hyperactivity disorder.⁵⁶ Similarly, primary care physicians now deliver many OB-GYN services. A 2023 study estimated that primary care physicians conducted 39% of preventive gynecological and women's health visits for women aged 18-44.⁵⁷ Women residing in rural areas, having a low socioeconomic status, and being over the age of 45 were more likely to see family medicine physicians or internists for OB-GYN services.^{58,59}

Chronic Health Conditions

Approximately 60% of adult Americans live with a chronic disease, with 40% of adult Americans having two or more chronic conditions.⁶⁰ With such a high percentage of the U.S. population living with chronic diseases, access to preventive care services, early detection, and regular management of chronic conditions are crucial. Experts acknowledge that episodic health care services, delivered in hospitals, are often insufficient in alleviating the impact of chronic disease on Americans' health. This suggests that primary care and a community-based approach are needed to ensure easy and affordable access to healthcare and improved patient outcomes.^{61,62,63,64,65}

Life Expectancy

People with better access to primary care live longer. A study of data from 2005-2015 found that an increase of 10 primary care physicians per 100,000 population was associated with a substantial increase in life expectancy (51.5 days), more than twice as large as the increase resulting from 10 additional specialist physicians per 100,000 population (19.2 days).⁶⁶ With an estimated average life expectancy at birth of 76.4 years, the U.S. ranks 32nd out of 38 Organization for Economic Cooperation and Development (OECD) countries (Table 7).⁶⁷ The U.S. also ranks lower in maternal mortality and infant mortality.

Table 7. Life Expectancy at Birth, Maternal Mortality and Infant Mortality in the U.S. and OECD Countries

Country	Life Expectancy at Birth ^a	Maternal Mortality ^b	Infant Mortality ^c
Australia	83.3	2.0	3.2
Canada	81.6	8.4	4.5
United Kingdom	80.4	N/A	3.8
Japan	84.5	2.7	1.8
United States	76.4	23.8	5.4
OECD countries average	80.3	10.5	4.0

Note. OECD Data Explorer (<https://data-explorer.oecd.org/>). ^a Life expectancy at birth in years in 2021, except Great Britain which is for 2020. ^b Deaths per 100,000 live births in 2020. ^c Infant deaths per 1,000 live births in 2020.

Conclusions

The importance of primary care cannot be overstated. Primary care is often the first contact a patient will have with the health care workforce and sets the trajectory for a positive or negative patient experience and outcome. A high-functioning primary care system treats illnesses and injuries before they become severe, provides ongoing care to mitigate chronic conditions, identifies when more specialized care is required, and connects the patient with a clinician. When primary care does not function as intended, patient issues can compound and become increasingly more difficult to treat and resolve.

The U.S. primary care system faces several challenges in the coming years. Barriers to health care access and shortages of providers result in uneven use of services. Because the primary care workforce is not distributed equally among geographic areas, many rural areas face low rates of physicians. Lower compensation compared to nonprimary care specialties and heightened stress and burnout (especially in the aftermath of COVID-19) are challenges in attracting and retaining new clinicians. The population of the United States will change in the future as will the methods to care for it.

Appendix

Table A. Ratios of Primary Care Physicians per 100,000 Population by U.S. State, 2022

State	Ratio	State	Ratio	State	Ratio
Alabama	68.9	Kentucky	70.2	North Dakota	85.8
Alaska	106.2	Louisiana	79.9	Ohio	85.0
Arizona	73.1	Maine	115.1	Oklahoma	64.4
Arkansas	76.2	Maryland	98.1	Oregon	103.9
California	89.7	Massachusetts	118.1	Pennsylvania	89.9
Colorado	90.2	Michigan	91.7	Rhode Island	110.4
Connecticut	95.3	Minnesota	97.9	South Carolina	74.0
Delaware	86.8	Mississippi	60.9	South Dakota	89.5
District of Columbia	177.2	Missouri	78.9	Tennessee	76.1
Florida	78.1	Montana	88.6	Texas	66.2
Georgia	72.7	Nebraska	84.8	Utah	60.2
Hawaii	103.4	Nevada	64.2	Vermont	123.0
Idaho	66.7	New Hampshire	94.5	Virginia	84.6
Illinois	91.0	New Jersey	85.0	Washington	93.2
Indiana	71.7	New Mexico	82.6	West Virginia	90.9
Iowa	77.9	New York	93.2	Wisconsin	88.5
Kansas	85.9	North Carolina	80.8	Wyoming	73.1

Note. 2022 AMA Physician Professional Data and U.S. Census Bureau estimates of State Population Totals for 2022 (<https://www.census.gov/data/tables/time-series/demo/popest/2020s-state-total.html>).

¹ Sullivan E.E., Etz R.S., Gonzalez M.M., Reves S.R., Deubel J., Stange K.C., Green L.A., Bitton A., Griffiths E.P., Sinsky C.A., & Linzer M. (2023). Primary care in peril: How clinicians view the problems and solutions. *NEJM Catalyst*. 4(6). <https://doi.org/10.1056/CAT.23.0029>

² Committee on the Future of Primary Care, Donaldson M., Yordy K., & Lohr K. (1996) Primary Care: America's Health in a New Era [online]. Washington, DC: National Academies Press (US). Retrieved October 24, 2023. <https://www.ncbi.nlm.nih.gov/books/NBK232631/>

³ Sharma G., Fletcher K.E., Zhang D., Kuo Y.F., Freeman J.L., & Goodwin J.S. (2009). Continuity of outpatient and inpatient care by primary care physicians for hospitalized older adults. *JAMA*. 301(16):1671-80. <https://doi.org/10.1001/jama.2009.517>

⁴ The American Academy of Family Physicians. Primary Care [online]. Leawood, Kansas: American Academy of Family Physicians. Retrieved on October 26, 2023. <https://www.aafp.org/about/policies/all/primary-care.html>

⁵ Jetty A., Petterson S., Westfall J.M., & Jabbarpour Y. (2021). Assessing primary care contributions to behavioral health: a cross-sectional study using medical expenditure panel survey. *Journal of Primary Care & Community Health*. 12(1). <https://doi.org/10.1177/21501327211023871>

⁶ Coffman M., Wilkinson E., & Jabbarpour Y. (2020). Despite adequate training, only half of family physicians provide women's health care services. *Journal of the American Board of Family Medicine*. 33(2):186-188. <https://doi.org/10.3122/jabfm.2020.02.190293>

⁷ Starfield B., Shi L., Macinko J. (2005). Contribution of primary care to health systems and health. *Milbank Quarterly*. 83(3): 457-502. <https://doi.org/10.1111/j.1468-0009.2005.00409.x>

⁸ Based on an analysis of the American Medical Association's 2022 AMA Physician Professional Data.

⁹ American Association of Medical Colleges. 2022 Physician specialty data report. Percentage change in the number of active physicians by specialty, 2016-2021. Retrieved October 30, 2023. <https://www.aamc.org/data-reports/workforce/data/percentage-change-number-active-physicians-specialty-2016-2021>

¹⁰ American Association of Nurse Practitioners. (2024, February). Workforce Survey- NP Fact Sheet. Retrieved April 12, 2024. <https://www.aanp.org/about/all-about-nps/np-fact-sheet>

- ¹¹ National Commission on Certification of Physician Assistants. (2024). NCCPA 2023 Annual Report: Statistical profile of certified PAs by specialty. Retrieved August 8, 2024. <https://www.nccpa.net/wp-content/uploads/2024/07/2023-Statistical-Profile-of-Board-Certified-PAs-by-Specialty-Annual-Report.pdf>
- ¹² United States Department of Labor, Bureau of Labor Statistics. Current Population Survey (CPS). 2023 annual data. Retrieved August 8, 2024. <https://www.bls.gov/cps/tables.htm>
- ¹³ Health Resources and Services Administration. Area Health Resources Files (AHRF) dashboard: 2018-2022 data from the American Community Survey. Retrieved August 30, 2024. <https://data.hrsa.gov/topics/health-workforce/ahrf>
- ¹⁴ Agency for Healthcare Research and Quality. Fact sheet: The distribution of the U.S. primary care workforce. Content last reviewed July 2018. Retrieved October 30, 2023. <https://www.ahrq.gov/research/findings/factsheets/primary/pcwork3/index.html>
- ¹⁵ Frazee T.K., Lewis V.A., Wood A., Newton H., & Colla C.H. (2022). Configuration and delivery of primary care in rural and urban settings. *Journal of General Internal Medicine*. 37(12):3045-3053. <https://doi.org/10.1007/s11606-022-07472-x>
- ¹⁶ Zhang D., Son H., Shen Y., Chen Z., Rajbhandari-Thapa J., Li Y., Eom H., Bu D., Mu L., Li G., & Pagán J.A. (2020). Assessment of changes in rural and urban primary care workforce in the United States from 2009 to 2017. *JAMA Network Open*. 3(10). <https://doi.org/10.1001/jamanetworkopen.2020.22914>
- ¹⁷ U.S. Census Bureau. Annual estimates of the resident population for counties: April 1, 2020 to July 1, 2023. Retrieved May 20, 2024. <https://www.census.gov/data/tables/time-series/demo/popest/2020s-counties-total.html>
- ¹⁸ American Association of Physician Assistants. (2022, September 15). Data brief: PA interest in rural locations, medically underserved areas, and health professional shortage areas. November 2021 Practice and 2022 AAPA Student Surveys. Retrieved October 30, 2023. <https://www.aapa.org/download/103451?tmstv=1685972579>
- ¹⁹ American Association of Physician Assistants. (2022). AAPA Data Dose. Physician assistants interest in rural practice? Retrieved October 30, 2023. <https://www.aapa.org/download/109210?tmstv=1685972332>
- ²⁰ Health Resources and Services Administration. (2024). Designated Health Professional Shortage Areas Statistics. Third Quarter of Fiscal Year 2024.
- ²¹ Health Resources and Services Administration's Workforce Projections. (2024, November). Retrieved November 7, 2024. <https://data.hrsa.gov/topics/health-workforce/workforce-projections>
- ²² Phillips R.L., Doodoo M.S., & Petterson S. (2009). Specialty and geographic distribution of the physician workforce: what influences medical student and resident choices? The Robert Graham Center. <https://www.graham-center.org/dam/rgc/documents/publications-reports/monographs-books/Specialty-geography-compressed.pdf>
- ²³ Nurse Journal Staff, *Nurse Journal*. (2022, July 18). Retrieved October 30, 2023. <https://nursejournal.org/careers/primary-care-nurse/salary/>
- ²⁴ Bureau of Labor Statistics. (2023, September 6). Occupational Outlook Handbook: Physician Assistants. Retrieved August 8, 2024. <https://www.bls.gov/ooh/healthcare/physician-assistants.htm>
- ²⁵ American Academy of Family Physicians. (2023). Position paper. Family physician burnout: Well-Being, and professional satisfaction. Retrieved October 31, 2023. <https://www.aafp.org/about/policies/all/family-physician-burnout.html>
- ²⁶ Eden A.R., Jabbarpour Y., Morgan Z.J., Wilkinson E., & Peterson L.E. (2020). Burnout among family physicians ... *Journal of the American Board of Family Medicine*. 33(3):355-356. <https://doi.org/10.3122/jabfm.2020.03.190319>
- ²⁷ Medscape. (2024). We have much work to do: Physician burnout & depression report. Retrieved August 20, 2024. <https://www.medscape.com/slideshow/2024-lifestyle-burnout-6016865#3>
- ²⁸ Shanafelt T.D., Hasan O., Dyrbye L.N., Sinsky C., Satele D., Sloan J., & West C.P. (2015). Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clinic Proceedings*. 90(12):1600-13. <https://doi.org/10.1016/j.mayocp.2015.08.023>
- ²⁹ Shanafelt T.D., West C.P., Sinsky C., Trockel M., Tutty M., Wang H., Carlasare L.E., & Dyrbye L.N. (2022). Changes in burnout and satisfaction with work-life integration in physicians and the general US working population between 2011 and 2020. *Mayo Clinic Proceedings*. 97(3):491-506. <https://doi.org/10.1016/j.mayocp.2021.11.021>
- ³⁰ Shanafelt T.D., West C.P., Dyrbye L.N., Trockel M., Tutty M., Wang H., Carlasare L.E., & Sinsky C. (2022) Changes in burnout and satisfaction with work-life integration in physicians during the first 2 years of the COVID-19 pandemic. *Mayo Clinic Proceedings*. 97(12):2248-2258. <https://doi.org/10.1016/j.mayocp.2022.09.002>
- ³¹ Patel R.S., Bachu R., Adikey A., Malik M., & Shah M. (2018). Factors related to physician burnout and its consequences: A Review. *Behavioral Science (Basel)*. 8(11):98. <https://doi.org/10.3390/bs8110098>
- ³² Pompeii L., Benavides E., Pop O., Rojas Y., Emery R., Delclos G., Markham C., Oluyomi A., Vellani K., & Levine N. (2020). Workplace violence in outpatient physician clinics: A systematic review. *International Journal of Environmental Research and Public Health*. 17(18):6587. <https://doi.org/10.3390/ijerph17186587>
- ³³ Grossman D.C. & Choucair B. (2019). Violence and the U.S. health care sector: Burden and response. *Health Affairs (Millwood)*. 38(10):1638-1645. <https://doi.org/10.1377/hlthaff.2019.00642>
- ³⁴ United States Department of Labor, Bureau of Labor Statistics. Injuries, Illnesses, and Fatalities. Retrieved October 31, 2023. <https://www.bls.gov/iif/home.htm>
- ³⁵ Patel S.Y., Mehrotra A., Huskamp H.A., Uscher-Pines L., Ganguli I., & Barnett M.L. (2021). Variation in telemedicine use and outpatient care during the COVID-19 pandemic in the United States. *Health Affairs (Millwood)*. 40(2):349-358. <https://doi.org/10.1377/hlthaff.2020.01786>

- ³⁶ Callaghan T., McCord C., Washburn D., Goidel K., Schmit C., Nuzhath T., Spiegelman A., & Scobee J. (2022). The changing nature of telehealth use by primary care physicians in the United States. *Journal of Primary Care & Community Health*. 13:1-9. <https://doi.org/10.1177/21501319221110418>
- ³⁷ Gajarawala S.N. & Pelkowski J.N. (2021). Telehealth benefits and barriers. *The Journal for Nurse Practitioners*. 17(2):218-221. <https://doi.org/10.1016/j.nurpra.2020.09.013>
- ³⁸ Department of Health and Human Services. Understanding telehealth. Retrieved October 31, 2023. <https://telehealth.hhs.gov/patients/understanding-telehealth>
- ³⁹ United States Census Bureau. Historical population change data (1910-2020). Retrieved October 31, 2023. <https://www.census.gov/data/tables/time-series/dec/popchange-data-text.html>
- ⁴⁰ United States Census Bureau. 2023 National Population Projections Tables: Main Series. Table 1. Projected Population and Components of Change. Retrieved August 9, 2024. <https://www2.census.gov/programs-surveys/popproj/tables/2023/2023-summary-tables/np2023-t1.xlsx>
- ⁴¹ United States Census Bureau. 2023 National Population Projections Tables: Main Series. Table 2. Projected Population by Age Group and Sex. Retrieved August 9, 2024. <https://www2.census.gov/programs-surveys/popproj/tables/2023/2023-summary-tables/np2023-t2.xlsx>
- ⁴² Willis J., Antono B., Bazemore A., Jetty A., Petterson S., George J., Rosario B.L., Scheufele E., Rajmane A., Dankwa-Mullan I., & Rhee K. (2020). The State of primary care in the United States: A Chartbook of facts and statistics. Retrieved October 31, 2023. <https://www.graham-center.org/content/dam/rgc/documents/publications-reports/reports/PrimaryCareChartbook2021.pdf>
- ⁴³ United States Census Bureau. 2023 National Population Projections Tables: Main Series. Table 3. Projected Population by Five-Year Age Group and Sex. Retrieved August 9, 2024. <https://www2.census.gov/programs-surveys/popproj/tables/2023/2023-summary-tables/np2023-t3.xlsx>
- ⁴⁴ Avery D.M.J., Bell J.G., Skinner C., & Geno C.E. (2018). Family physicians providing rural obstetric care makes good business sense. *Clinical Obstetrics, Gynecology, and Reproductive Medicine*. 4(3):1-3. <https://doi.org/10.15761/COGRM.1000219>
- ⁴⁵ Eden A.R., Barreto T., & Hansen E.R. (2019). Experiences of new family physicians finding jobs with obstetrical care in the USA. *Family Medicine and Community Health*. 7(3). <https://doi.org/10.1136/fmch-2018-000063>
- ⁴⁶ United States Government Accountability Office. (2022). Maternal health: Availability of hospital-based obstetric care in rural areas. Retrieved October 31, 2023. <https://www.gao.gov/products/gao-23-105515>
- ⁴⁷ United States Census Bureau. 2023 National Population Projections Tables: Main Series. Table 4. Projected Population by Sex, Race, and Hispanic Origin. Retrieved August 9, 2024. <https://www2.census.gov/programs-surveys/popproj/tables/2023/2023-summary-tables/np2023-t4.xlsx>
- ⁴⁸ Basu S., Berkowitz S.A., Phillips R.L., Bitton A., Landon B.E., & Phillips R.S. (2019). Association of primary care physician supply with population mortality in the United States, 2005-2015. *JAMA Internal Medicine*. 179(4):506-514. <https://doi.org/10.1001/jamainternmed.2018.7624>
- ⁴⁹ Department of Human and Health Services, Office of Disease Prevention and Health Promotion. Healthy People 2030. Access to primary care. Retrieved October 31, 2023. <https://health.gov/healthypeople/priority-areas/social-determinants-health/literature-summaries/access-primary-care>
- ⁵⁰ Douthit N., Kiv S., Dwolatzky T., Biswas S. (2015). Exposing some important barriers to health care access in the rural USA. *Public Health*. 129(6):611-620. <https://doi.org/10.1016/j.puhe.2015.04.001>
- ⁵¹ Huffstetler A.N., Jetty A., Greiner A., & Jabbarpour Y. (2023). Relationships matter: Primary care physicians and usual sources of care. *American Family Physician*. 107(4):356-357.
- ⁵² Lee D.C., Shi L., Wang J., & Sun G. (2023). Usual source of care and access to care in the US: 2005 vs. 2015. *PLoS One*. 18(1). <https://doi.org/10.1371/journal.pone.0278015>
- ⁵³ Agency for Health Research and Quality. MEPS Topics: Usual Source of Care. Retrieved October 31, 2023. https://meps.ahrq.gov/mepsweb/data_stats/MEPS_topics.jsp?topicid=44Z-1
- ⁵⁴ Young R.A. & Sundermeyer R.L. (2018). Family medicine and obstetrics: Let's stop pretending. *Journal of the American Board of Family Medicine*. 31(3):328-331. <https://doi.org/10.3122/jabfm.2018.03.180087>
- ⁵⁵ Rotenstein L.S., Edwards S.T., & Landon B.E. (2023). Adult primary care physician visits increasingly address mental health concerns. *Health Affairs (Millwood)*. 42(2):163-171. <https://doi.org/10.1377/hlthaff.2022.00705>
- ⁵⁶ Reed S.J., Shore K.K., & Tice J.A. (2016). Effectiveness and value of integrating behavioral health into primary care. *JAMA Internal Medicine*. 176(5):691-692. <https://doi.org/10.1001/jamainternmed.2016.0804>
- ⁵⁷ Attanasio L., Ranchoff B., Jeung C., Goff S., & Geissler K. (2023). Preventive care visits with OB/GYNs and generalist physicians among reproductive-age women with chronic conditions. *Health Services Research*. 58(1):207-215. <https://doi.org/10.1111/1475-6773.14100>
- ⁵⁸ Raffoul M.C., Petterson S.M., Rayburn W.F., Wingrove P., & Bazemore A.W. (2016). Office visits for women aged 45-64 years according to physician specialties. *Journal of Women's Health (Larchmt)*. 25(12):1231-1236. <https://doi.org/10.1089/jwh.2015.5599>

-
- ⁵⁹ Simon A.E. & Uddin S.F.G. (2017). Trends in seeing an obstetrician-gynecologist compared with a general physician among U.S. women, 2000-2015. *Obstetrics & Gynecology*. 130(4):677-683. <https://doi.org/10.1097/AOG.0000000000002248>
- ⁶⁰ Buttorff C., Ruder T., & Bauman M. (2017). Multiple Chronic Conditions in the United States. RAND Corporation. <https://www.rand.org/pubs/tools/TL221.html>
- ⁶¹ Decker S.L., Schappert S.M., & Sisk J.E. (2009). Use of medical care for chronic conditions. *Health Affairs (Millwood)*. 28(1):26-35. <https://doi.org/10.1377/hlthaff.28.1.26>
- ⁶² Reynolds R., Dennis S., Hasan I., Slewa J., Chen W., Tian D., Bobba S., & Zwar N. (2018). A systematic review of chronic disease management interventions in primary care. *BMC Family Practice*. 19(1):11. <https://doi.org/10.1186/s12875-017-0692-3>
- ⁶³ Smith S.M., Wallace E., O'Dowd T., & Fortin M. (2021). Interventions for improving outcomes in patients with multimorbidity in primary care and community settings. *Cochrane Database of Systematic Reviews*. 1(1). <https://doi.org/10.1002/14651858.CD006560.pub4>
- ⁶⁴ Gulliford M. (2017). Access to primary care and public health. *The Lancet Public Health*. 2(12):e532-e533. [https://doi.org/10.1016/S2468-2667\(17\)30218-9](https://doi.org/10.1016/S2468-2667(17)30218-9)
- ⁶⁵ Schneider E.C., Shah A., Doty M.M., Tikkanen R., Fields K., & Williams R.D. (2021). Mirror, mirror 2021: Reflecting poorly. The Commonwealth Fund. Retrieved October 31, 2023. <https://www.commonwealthfund.org/publications/fund-reports/2021/aug/mirror-mirror-2021-reflecting-poorly>
- ⁶⁶ Basu S., Berkowitz S.A., Phillips R.L., Bitton A., Landon B.E., & Phillips R.S. (2019). Association of primary care physician supply with population mortality in the United States, 2005-2015. *JAMA Internal Medicine*. 179(4):506-514. <https://doi.org/10.1001/jamainternmed.2018.7624>
- ⁶⁷ American Public Health Associations and America's Health Ranking. (2023). America's Health Rankings 2023 Annual Report. https://assets.americashealthrankings.org/app/uploads/ahr_2023annual_comprehensivereport_final2-web.pdf